

University of Groningen

## Fusional vergence eye movements in microstrabismus

Boonstra, Frouke Nienke

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

1997

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Boonstra, F. N. (1997). *Fusional vergence eye movements in microstrabismus*. s.n.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## 2. Vergence and vergence eye movements

### 2.1 Introduction

Eye movements can be classified in various ways. Carpenter (1977) divided eye movements into catching or fast movements, such as saccades and the quick phase of the nystagmus, and holding or slow movements, such as vestibular movements, smooth pursuit and vergence.

Another division is between conjugate and disjunctive movements. Saccades, for example, are conjugate movements and vergence movements are disjunctive movements. Below we shall focus on disjunctive movements (vergence). See figure 2.1.



**figure 2.1**  
Classification of eye movements.

Throughout this thesis, vergence refers to the hypothesized underlying neuronal process. Convergence and divergence refer to the actual eye movements.

In 1893, Maddox divided horizontal vergence in four different types:

- accommodative vergence;
- fusional vergence;
- proximal vergence;
- tonic vergence.

An alternative classification of vergence was given by Crone (1979), who divided horizontal vergence into three mechanisms:

1. motor fusion (the F system);
2. synkinesis of near fixation, accommodation and miosis (the S system);
3. the slow mechanism of orthophorization, the O system

See also section 2.5.

In recent years, several authors have reported vergence eye movements which were faster than the fusional or accommodative vergence eye movements recorded in the usual classical experiments, such as:

- vergence mediated by saccades (Enright, 1984);
- rapid vergences (Erkelens, 1986; 1991);
- proximal vergence (Wick and Bedell, 1989);
- vergence saccades (Hyson, 1983).

These vergence movements have in common that they were recorded in more complex (natural) viewing conditions.

In this study the division of vergence types is based on the way the movement can be evoked. Therefore the different types of vergence are described according to the following classification:

- by disparity (section 2.2);
- by blur (section 2.3);
- in a complex stimulus environment (section 2.4).

## **2.2 Disparity Vergence**

### **2.2.1 Fusional vergence**

#### **2.2.1.1 Normal fusional vergence**

Fusion consists of both motor fusion and sensory fusion. The movement that serves to bring the images on corresponding retinal points is called motor fusion. With the following experiment motor fusion, or fusional vergence, is described (figure 2.1). A subject is binocularly fixating a visual target. A weak horizontal prism is suddenly placed in front of one eye; the right eye in this case (see figure 2.1<sup>a</sup>). The prism causes a shift of the image, which results in disparity. This disparity leads to a vergence movement (see figure 2.1b and 2.1c from one of our experiments). The subject is generally unaware of this (Carpenter, 1977).




figure 2.1<sup>a</sup>




figure 2.1<sup>b</sup>

In this example, just after the beginning of the vergence movement, a fast versional movement, a saccade, takes place. The saccade is evoked by the process of intermediate localisation, caused by the displacement of the prism in front of only one eye. After the saccade the vergence movement becomes more obvious (see figure 2.1<sup>c</sup>). By this (fusional) vergence movement, the image is again projected on corresponding areas in the retinae of both eyes (areas with a comparable localisation in space). The stimulus for this fusional vergence is disparity; therefore, the term disparity vergence will often be used instead of fusional vergence. Such a movement is illustrated in figure 2.1<sup>c</sup>.

